

Direct observations of the effects of aerosol loading on net ecosystem CO₂ exchanges over different landscapes

Dev Niyogi,¹ Hsin-I Chang,¹ V. K. Saxena,¹ Teddy Holt,^{1,2} Kiran Alapaty,³ Fitzgerald Booker,⁴ Fei Chen,⁵ Kenneth J. Davis,⁶ Brent Holben,⁷ Toshihisa Matsui,⁸ Tilden Meyers,⁹ Walter C. Oechel,¹⁰ Roger A. Pielke Sr.,⁸ Randy Wells,¹¹ Kell Wilson,⁹ and Yongkang Xue¹²

Received 5 July 2004; revised 2 September 2004; accepted 29 September 2004; published 29 October 2004.

[1] We present the first direct, multisite observations in support of the hypothesis that atmospheric aerosols affect the regional terrestrial carbon cycle. The daytime growing season (summer) CO₂ flux observations from six sites (forest, grasslands, and croplands) with collocated aerosol and surface radiation measurements were analyzed for high and low diffuse radiation; effect of cloud cover; and effect of high and low aerosol optical depths (AOD). Results indicate that, aerosols exert a significant impact on net CO₂ exchange, and their effect may be even more significant than that due to clouds. The response appears to be a general feature irrespective of the landscape and photosynthetic pathway. The CO₂ sink increased with aerosol loading for forest and crop lands, and decreased for grassland. The cause for the difference in response between vegetation types is hypothesized to be canopy architecture. **INDEX TERMS:** 0305 Atmospheric Composition and Structure: Aerosols and particles (0345, 4801); 0315 Atmospheric Composition and Structure: Biosphere/atmosphere interactions; 0345 Atmospheric Composition and Structure: Pollution—urban and regional (0305); 1610 Global Change: Atmosphere (0315, 0325); 1615 Global Change: Biogeochemical processes (4805). **Citation:** Niyogi, D., et al. (2004), Direct observations of the effects of aerosol loading on net ecosystem CO₂ exchanges over different landscapes, *Geophys. Res. Lett.*, 31, L20506, doi:10.1029/2004GL020915.

1. Introduction

[2] Photosynthesis removes large amounts of CO₂ from the atmosphere. Net global terrestrial carbon exchange was nearly neutral in the 1980's, but resulted in a carbon sink in the 1990's [Schimel et al., 2001]. CO₂ fertilization, land cover/land-use change, nitrogen loading, forest fires, and the regional hydrological cycle are some of the known factors affecting the carbon cycle [Nemani et al., 2002]. Recent studies suggest that clouds and aerosols released in

the atmosphere due to volcanic eruptions could also be important factors [Gu et al., 2003; Farquhar and Roderick, 2003; Krakauer and Randerson, 2003].

[3] Given that previous studies cite significant events such as volcanic eruptions as the cause for variability in the carbon cycle, and that the mechanisms responsible for modified photosynthetic rates are modulated by aerosol loading, we ask the question: *can we detect the effect of relatively routine aerosol variability on field measurements of CO₂ fluxes, and if so, how does the variability in aerosol loading affect CO₂ fluxes over different landscapes?* Further, since studies such as Krakauer and Randerson [2003] question the positive effects of aerosols on the terrestrial carbon cycle; and modeling analysis of Cohan et al. [2002] indicated that the aerosol effects on CO₂ fluxes could depend on cloudiness, we seek to find: *whether or not the direct observations indicate an increase or a decrease in field scale CO₂ fluxes?* Thus, even though the effects of clouds on CO₂ fluxes are well documented [Hollinger et al., 1998; Gu et al., 2002], studies linking direct observations of aerosol loading on surface CO₂ fluxes are lacking. Using field measurements, we present additional evidence of the importance of aerosol feedback on regional climate via the biogeochemical pathways affecting the terrestrial carbon cycle.

2. Data and Methods

[4] We used CO₂ flux (*F_c*) data from the AmeriFlux network [Baldocchi et al., 2001], and cloud-free aerosol optical depth (AOD) data from NASA Aerosol Robotic Network: AERONET [Holben et al., 2001] for assessing the effect of aerosol loading on net ecosystem exchange (NEE). Six locations had concurrent *F_c* and AOD observations. The landscapes (locations and period with concurrent AOD and *F_c* data) were: broadleaf deciduous forest (Walker Branch, TN, 2000), mixed forest (Willow Creek/Lost Creek, WI, 2000–01), crops (winter wheat: Ponca, OK, 1998–99; alternate soybean or corn: Bondville, IL, 1998–2002), and grassland (Barrow, AK, 1999; Shidler, OK, 1998–99).

¹Department of MEAS, North Carolina State University, Raleigh, North Carolina, USA.

²Also at Naval Research Laboratory, Monterey, CA, USA.

³University of North Carolina at Chapel Hill, Raleigh, North Carolina, USA.

⁴USDA-ARS Air Quality–Plant Growth and Development Unit, Raleigh, North Carolina, USA.

⁵National Center for Atmospheric Research, Boulder, Colorado, USA.

⁶Department of Meteorology, Pennsylvania State University, University Park, Pennsylvania, USA.

⁷NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.

⁸Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado, USA.

⁹Atmospheric Turbulence and Diffusion Division, National Oceanic and Atmospheric Administration, Oak Ridge, Tennessee, USA.

¹⁰San Diego State University, San Diego, California, USA.

¹¹Department of Crop Science, North Carolina State University, Raleigh, North Carolina, USA.

¹²Department of Geography, University of California, Los Angeles, Los Angeles, California, USA.




Figure 1a. Observed 30-minute averaged daytime observations of CO₂ flux and global irradiance for 1996–2000 summer (June–August). 2nd polynomial best fit ($n = 3177$, $p < 0.05$) are shown. Solid line: high diffuse regimes ($Rd/Rg > 0.6$, $r = 0.78$); dashed line: low diffuse radiation regime ($Rd/Rg < 0.4$, $r = 0.67$). See color version of this figure in the HTML.

[5] All data were quality assured by graphical and statistical means. Periods with either F_c or AOD measurements missing were eliminated. Daytime observations (since solar radiative effects will be studied) from June through August were selected. This period corresponds to the growing season, and includes the peak photosynthetic activity and capacity of the canopy.

[6] Data were clustered into three pairs to test the sensitivity of the F_c to the diffuse radiative flux fraction (DRF), and then to assess the impact of clouds as well as aerosol loading on the F_c .

[7] In analysis I, all observations, regardless of cloud cover or aerosol loading were clustered according to DRF [calculated as ratio of diffuse (Rd) to global irradiance (Rg)]. Data with $Rd/Rg > 0.6$ were labeled high diffuse regime, and those with $Rd/Rg < 0.4$ were labeled as a low diffuse regime. For analysis II, data were clustered according to cloudy and non-cloudy sky conditions. This was determined by analyzing the global irradiance time series plots, GOES visible cloud images, and weather reports for each day [Gu *et al.*, 1999]. In analysis III, observations were analyzed according to the AOD values, for clear sky (i.e., no clouds) conditions.

3. Results and Discussion

[8] Since similar analysis was conducted for each site, we will discuss the results for one site (Walker Branch) in detail and summarize the results for all the sites.

3.1. Effect of Diffuse Radiation

[9] Figure 1a shows the observed daytime F_c (30-min averages) for high and low DRF clusters (i.e., Analysis I) for the 1996 to 2000 summer (June–August). The F_c increase in magnitude (in the figures a negative value indicates a net flux into the vegetation, i.e., a sink) as a function of Rg. Additionally, for the same irradiance, the surface F_c increases (in magnitude) with increasing DRF. For example, for $Rg = 500 \text{ Wm}^{-2}$, the F_c is $13 \mu\text{molm}^{-2}\text{s}^{-1}$ for low DRF, and $20 \mu\text{molm}^{-2}\text{s}^{-1}$ i.e., about 50% higher, for the high DRF. Observations for the




Figure 1b. Normalized daily CO₂ fluxes and diffuse fractions from the period shown in Figure 1a.

three summer months over five years clearly indicate a significant increase in daytime NEE for larger DRF for similar global irradiance.

[10] The data from 1000 to 1600 LT, shown in Figure 1a were further averaged to yield a ‘daily’ value. The early morning and evening period were eliminated to avoid confounding due to low solar angles on high DRF caused independent of cloud cover or aerosol loading [Gu *et al.*, 1999]. The ‘daily’ averaged data are plotted in Figure 1b, with the F_c values normalized by Rg. A linear relation is obtained between higher DRF and the F_c values (both normalized for Rg). The best fit ($n = 178$, $r = 0.75$, $p < 0.05$) indicates that higher DRF enhances photosynthetic fluxes by about 30% at this study site.

3.2. Effect of Clouds

[11] The effect of increased mid-day DRF on F_c values (Figures 1a–1b) can be related to increased cloud cover and/or aerosol loading. Typically, for cloudy, overcast conditions the DRF is close to one. Hence the F_c data were clustered into ‘clear’ (i.e., non-cloudy) and ‘cloudy’ regimes (i.e., Analysis II, Figure 2a). The results are similar to those obtained in prior studies [Hollinger *et al.*, 1994; Gu *et al.*, 1999; Roderick *et al.*, 2001]. That is, under cloudy conditions the F_c values are larger for similar Rg. As in




Figure 2a. Effect of cloudiness on CO₂ flux. Solid line: ‘Cloudy’ sky ($r = 0.75$, $n = 1278$); dashed line: ‘Clear’ (non-cloudy) sky ($r = 0.74$, $n = 1395$). See color version of this figure in the HTML.




Figure 2b. Normalized 30-minute averaged CO_2 flux and diffuse radiation fraction (DRF) clustered for cloudy (+) and non-cloudy conditions. Solid line is best fit ($r = 0.14$, $n = 2479$). See color version of this figure in the HTML.

Analysis I, the 30-minute F_c values were normalized by R_g (Figure 2b). The normalized F_c show two distinct clusters for cloudy and clear (non-cloudy) sky conditions. CO_2 fluxes under cloudy conditions (i.e., high DRF) are typically larger than those under non-cloudy conditions with the same R_g (i.e., lower DRF).

3.3. Effect of Aerosol Loading

[12] As seen in Figures 2a–2b, some cloud-free days also have relatively high DRF values. This can be due to aerosols. Hence, the clear sky conditions data were analyzed further as a function of aerosol loading (through aerosol optical depth: AOD). For the Walker Branch site, coincident F_c and AOD data were available for June and July 2000 (Figure 3).

[13] The results indicate that the CO_2 flux is typically higher for larger aerosol loading. Another noteworthy feature of Figure 3 is that even under high aerosol loading, the R_g can be high ($\sim 900 \text{ W m}^{-2}$).

[14] The increase in F_c with aerosol loading is likely to be the result of larger DRF. Consequently, the effect of AOD on DRF was studied. DRF shows a near linear relation with aerosol loading ($r = 0.92$, $n = 119$, not shown). Therefore, the variation in CO_2 fluxes can be considered to be an indirect




Figure 3. 30-minute averaged daytime observations of CO_2 flux and global irradiance during June–July of 2000. 2nd polynomial best fit is shown ($n = 255$), solid: $\text{AOD} > 0.6$ ($r = 0.81$); dashed: $\text{AOD} < 0.4$ ($r = 0.72$).




Figure 4. Aerosol optical depth (AOD) and CO_2 flux at the Walker Branch site for June–July 2000. With increasing aerosol loading, the landscape is a larger CO_2 sink. Solid line is best fit ($n = 43$, $r = 0.76$). See color version of this figure in the HTML.

effect of increased regional aerosol loading. Indeed, a nearly linear relation is obtained between AOD and the F_c under the clear sky conditions (Figure 4). With increasing AOD, the surface CO_2 fluxes are consistently larger.

3.4. Comparing the Effect of Aerosol Loading on CO_2 Fluxes Over Different Landscapes

[15] To investigate the regional effect of aerosol loading on field-scale NEE, the analysis described for the Walker Branch forest site was repeated for the five other sites (Figure 5). The sites represented the following landscapes (and photosynthetic pathways): winter wheat (C3), corn (C4), soybean (C3), grasslands (C3/C4), and mixed forest (C3).

[16] For all the sites, field-scale NEE varies with AOD, and every landscape has a different response. The woody (Figure 5a) and agricultural (Figure 5b) sites show an increase in the field-scale CO_2 flux ‘sink’ due to aerosol loading. Interestingly, both the grassland sites (Figure 5c) show an opposite response, and indicate a decreased field-scale CO_2 flux ‘sink’ with increased aerosol loading. Reviewing the best fits, the effect of aerosol loading on F_c is largest for C4 grassland (Shidler), and crops (corn; Bondville 1999, 2001), and relatively less on C3 crops (winter wheat, Ponca; and soybean, Bondville 1998, 2000, 2002). The F_c measurements over trees (forest sites) are also sensitive to the aerosol loading. Thus, both the canopy architecture (and hence the canopy scale radiative feedback on photosynthesis) and the photosynthesis pathway appear to be important factors. Additionally, there is significant scatter in the relationship between AOD and F_c , indicating other environmental variables (beyond aerosol loading) also influence the results. Indeed for all the landscapes, those variables that are known to affect photosynthesis rates (such as leaf area index and soil moisture availability) were also found to be significant in modulating the CO_2 fluxes (not shown). The results, however, clearly indicate that aerosol loading has a significant impact on the net ecosystem CO_2 exchange over terrestrial landscapes.

[17] For the results discussed above (Figure 5) 500 nm AOD data were chosen since it corresponded to a PAR wavelength. The AERONET AODs are centered over seven wavelengths (340 nm to 1020 nm), and the analysis was extended for all AOD wavelengths.




Figure 5. Relation between aerosol optical depth (AOD) and CO₂ flux at different sites and landscapes. The best fits and the periods for which data were used are also shown. (a) Trees- WB: Walker Branch ($r = 0.76$, $n = 43$); WC,LC: Willow Creek, Lost Creek ($r = 0.52$, $n = 124$); (b) Crops - BV: Bondville ($r = 0.44$, $n = 150$, for BV(98,00,02); $n = 82$, $r = 0.61$ for BV(99, 01); and (c) Grass- Barrow ($r = 0.39$, $n = 23$) and Shidler ($r = 0.55$, $n = 124$). Aerosols can increase (decrease) CO₂ fluxes/sink potential over forest and croplands (grasslands). C4 vegetation [Shidler; BV(99,01)] show largest sensitivity, C3 crops/grasslands show least, and trees show a moderately high influence of aerosols on CO₂ fluxes. See color version of this figure in the HTML.

[18] The AOD – CO₂ flux relation is sensitive to the choice of wavelength used and different landscapes may show sensitivity to different wavelengths. For example, the slope of the CO₂ flux – AOD best fit varies between -1.4 (for 340 nm) to -1.54 (for 1020 nm) radians for the deciduous forest site, and corresponding variation is comparatively less (from -1.53 to -1.55 radians) for the cropland. Thus NEE over a woody landscape could be even more sensitive to the aerosol loading than discussed in the analysis above. However, this does not alter the conclusion that aerosol loading can influence the terrestrial CO₂ fluxes.

4. Conclusions

[19] Study results suggest that aerosol induced radiative effect is an important modulator of regional carbon cycles. For the different study sites, DRF affected CO₂ fluxes, with an increase in DRF correlating with higher CO₂ flux values (sink) for trees and crops; and a lower sink for grasslands. The effect was clearly seen under cloudy conditions, during which the DRF was close to unity. It was also identified under higher aerosol loading in non-cloudy sky conditions. Aerosols can therefore routinely influence surface irradiance and hence the terrestrial CO₂ flux and regional carbon cycle.

[20] The reason for increased CO₂ fluxes with increasing DRF for forests and croplands is considered to result from an increase in the vegetative canopy fraction that is receiving illumination (without photosaturation). The advantage of increased diffuse radiation under clear sky, high aerosol conditions does not appear to be available to grasslands due to the canopy architecture. Additional confounding effects due to temperature, and humidity might exist and should be explored. Data from the early morning and evening period will also have high DRF but were eliminated in our analysis. For the whole-day carbon exchange, other factors that alter plant response may offset the effects of aerosol loading.

[21] Aerosols are abundant in the environment and their effects on climate are only poorly understood. Aerosols can alter irradiance at the top of atmosphere and even more profoundly at the surface and affect the biosphere [Schwartz,

1996]. Changes in DRF, due to aerosol loading, appear to have the potential to alter the terrestrial carbon exchange.

[22] Past studies on the impact of diffuse radiation on CO₂ flux focused on either the effect of cloudiness [e.g., Hollinger *et al.*, 1998] or the impact of volcanic aerosols [Gu *et al.*, 2003] on CO₂ flux exchange. Gu *et al.* [2002] analyzed field measurements across the United States and showed that NEE was larger for a higher diffuse fraction of the incoming radiation. Thus, even though field and model studies provide increasing evidence that photosynthesis rates and NEE will increase with DRF, the majority of these studies have been based on either episodic analysis of aerosol loading (i.e., effect of the Mt. Pinatubo eruption) or as the effect of cloudiness (which are conditions of high DRF). Thus our study is the first multi-site, observational analysis to investigate effects of persistent regional aerosol loading (which typically has a lifetime on the order of a week) on field NEE. Such an analysis is important for several reasons. First, the results provide evidence that routine aerosol loading due to natural or anthropogenic sources have the potential to influence regional CO₂ flux. Second, even though past studies indicate that for a given irradiance, NEE would be higher under cloudy conditions; cloudiness in itself may not be the dominant factor that increases the ability of a region to be a carbon sink. This is because even though it has a large DRF, total radiation is dramatically reduced. Hence cloudiness could even lead to lower, rather than higher, NEE over the region [cf. Krakauer and Randerson, 2003]. Alternatively, our results indicate that increasing aerosol loading will increase the diffuse fraction of the radiation without significantly reducing the total radiation itself and could be a prominent forcing affecting the CO₂ flux variability over a region. Thus, the potential of the vegetated land surface to be a sink for atmospheric carbon could depend on regional aerosol loading.

[23] **Acknowledgments.** Study benefited from support through ONR, NASA- Radiation Science (NNG04GB87G Drs. D. Anderson and T. Lee), as well as NASA- THP (NNG04GI84G Dr. J. Entin), NASA-IDS (NNG04GL61G Drs. J. Entin and G. Gutman), NSF (ATM-0233780,

Dr. S. Nelson), NCSU FRPD Award, NCAR Water Cycle Initiative, and NSF-AWARE (Dr. B. Kuo). Many thanks to Drs. Lianhong Gu and Gregg Marland, ORNL; Prof. James Randerson, UC Irvine for review and comments on the manuscript, and to Profs. S.B. Verma, UNL; D. Baldocchi, UC-Berkeley; P. Bolstad, U of MN.; for data/suggestions.

References

- Baldocchi, D., et al. (2001), FLUXNET: A new tool to study the temporal and spatial variability of ecosystem-scale carbon-dioxide, water vapor, and energy flux densities, *Bull. Am. Meteorol. Soc.*, *82*, 2415–2434.
- Cohan, D. S., et al. (2002), Impact of atmospheric aerosol light scattering and absorption on terrestrial net primary productivity, *Global Biogeochem. Cycles*, *16*(4), 1090, doi:10.1029/2001GB001441.
- Farquhar, G., and M. L. Roderick (2003), Pinatubo, diffuse light, and the carbon cycle, *Science*, *299*, 1997–1998.
- Gu, L., et al. (1999), Responses of net ecosystem exchanges of carbon dioxide to changes in cloudiness: Results from two North American deciduous forests, *J. Geophys. Res.*, *104*, 31,421–31,434.
- Gu, L., et al. (2002), Advantages of diffuse radiation for terrestrial ecosystem productivity, *J. Geophys. Res.*, *107*(D6), 4050, doi:10.1029/2001JD001242.
- Gu, L., et al. (2003), Response of a deciduous forest to the Mount Pinatubo eruption: Enhanced photosynthesis, *Science*, *299*, 2035–2038.
- Holben, B. N., et al. (2001), An emerging ground-based aerosol climatology: Aerosol Optical Depth from AERONET, *J. Geophys. Res.*, *106*, 12,067–12,097.
- Hollinger, D. Y., et al. (1998), Forest-atmosphere carbon dioxide exchange in eastern Siberia, *Agric. For. Meteorol.*, *90*, 291–306.
- Krakauer, N. Y., and J. T. Randerson (2003), Do volcanic eruptions enhance or diminish net primary production? Evidence from tree rings, *Global Biogeochem. Cycles*, *17*(4), 1118, doi:10.1029/2003GB002076.
- Nemani, R., et al. (2002), Recent trends in hydrologic balance have enhanced the terrestrial carbon sink in the United States, *Geophys. Res. Lett.*, *29*(10), 1468, doi:10.1029/2002GL014867.
- Roderick, M. L., et al. (2001), On the direct effect of clouds and atmospheric particles on the productivity and structure of vegetation, *Oecologia*, *129*, 21–30.
- Schimel, D. S., et al. (2001), Recent patterns and mechanisms of carbon exchange by terrestrial ecosystems, *Nature*, *414*, 169–172.
- Schwartz, S. E. (1996), The Whitehouse Effect—Shortwave radiative forcing of climate by anthropogenic aerosols: An overview, *J. Atmos. Sci.*, *27*, 359–382.
- K. Alapaty, University of North Carolina at Chapel Hill, Raleigh, NC 27695-8208, USA.
- F. Booker, USDA-ARS Air Quality–Plant Growth and Development Unit, Raleigh, NC 27695-8208, USA.
- H. Chang, T. Holt, D. Niyogi, and V. K. Saxena, Department of MEAS, North Carolina State University, Raleigh, NC 27695-8208, USA. (dev_niyogi@ncsu.edu)
- F. Chen, National Center for Atmospheric Research, Boulder, CO 80305, USA.
- K. J. Davis, Department of Meteorology, Pennsylvania State University, University Park, PA 16802, USA.
- B. Holben, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA.
- T. Matsui and R. A. Pielke Sr., Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523, USA.
- T. Meyers and K. Wilson, Atmospheric Turbulence and Diffusion Division, National Oceanic and Atmospheric Administration, Oak Ridge, TN 37830, USA.
- W. C. Oechel, San Diego State University, San Diego, CA 92182, USA.
- R. Wells, Department of Crop Science, North Carolina State University, Raleigh, NC 27695-8208, USA.
- Y. Xue, Department of Geography, University of California, Los Angeles, Los Angeles, CA 90095, USA.